

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 1 Jan 98	3. REPORT TYPE AND DATES COVERED Final Report 8/1/93 - 7/31/96	
4. TITLE AND SUBTITLE Molecular Approaches to Optical Biosensors			5. FUNDING NUMBERS N00014-93-1-1245	
6. AUTHOR(S) Carol A. Fierke				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Biochemistry Department, Box 3711 Duke University Medical Center Durham, NC 27710			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quicy St. Arlington, VA 22217-5000			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Distribution Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  The goal of this proposal was to develop methodologies for the optimization of field-deployable optical biosensors, in general, and, in particular, to optimize a carbonic anhydrase-based fiber optic zinc biosensor. We used both site-directed and random mutagenesis coupled with phage display to prepare and identify carbonic anhydrase (CA) variants that: 1) enhance the kinetics of metal $\geq 1000$ -fold such that zinc equilibration in the nM range occurs within seconds; 2) alter the zinc affinity by a factor of $10^7$ -fold (20 fM - 0.2 $\mu$ M) for use in a biosensor array; and 3) improve the metal detection limit by altering the transduction scheme. Additionally, we have developed phage display technology to screen for CA variants with alter zinc affinity.				
14. SUBJECT TERMS Metals, biosensors, phage display, mutagenesis			15. NUMBER OF PAGES 4	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

19990701 037

FINAL REPORT

Grant#: N00014-93-1-1245

PRINCIPAL INVESTIGATOR: Dr. Carol Ann Fierke

INSTITUTION: Duke University Medical Center

E-MAIL: fierke@biochem.duke.edu

GRANT TITLE: Molecular Approaches to Optical Biosensors

AWARD PERIOD: 1 August 1993 - 31 July 1996

OBJECTIVE: To develop methodologies for the optimization of field-deployable optical biosensors; to improve the zinc detection limit and speed of equilibration of a carbonic anhydrase-based fiber optic zinc biosensor.

APPROACH: Use both site-directed and random mutagenesis coupled with activity screens to prepare and identify carbonic anhydrase (CA) variants that: 1) increase the kinetics of metal binding; 2) alter the affinity and specificity of the metal site; and 3) improve the metal detection limit by altering the transduction scheme.

ACCOMPLISHMENTS:

*Site-directed mutagenesis*: To investigate the molecular mechanism for slow zinc equilibration we have prepared CA variants in two amino acids, Gln92 and Glu117, which form hydrogen bonds with His94 and His119. Substitution of Gln92 causes modest increases in both the zinc  $K_D$  and the dissociation rate constant. Substitution of Glu117 also causes modest increases (<10-fold) in the zinc  $K_D$ ; however, the **zinc association and dissociation rate constants increase dramatically** (1000-fold) with little apparent effect on catalytic activity (except for Glu117→Gln) or protein stability. In fact, the Glu117→Gln variant decreases the half-time for equilibration with 0.5 nM zinc from 1 hour to 5 seconds.

We have demonstrated that the histidines that coordinate zinc in CAII can be functionally replaced by asparagine, glutamine, glutamate, aspartate, and cysteine. This causes large ( $10^5$ -fold) increases in the zinc dissociation rate constant which are accompanied by large decreases in the binding constants for both zinc. These variants may be particularly useful for detecting nM concentrations of zinc in an equilibrium manner. We have also inserted a fourth protein ligand into the zinc coordination polyhedron of carbonic anhydrase II which **increases the metal affinity 130-fold to 20 fM**. To date, we have prepared a collection of CAII variants with zinc affinity varying

by a factor of  $10^7$ -fold (20 fM - 0.2  $\mu$ M) for use in a biosensor array.

Additionally, we have supplied our collaborator, Dr. Richard Thompson, CA variants containing one cysteine substituted at a number of positions at different distances from zinc, including Y7C (12Å), N67C (9.6Å), H64C (9Å), V143C (7.1Å) and L198C (7.8Å) CAII. Dr. Thompson's laboratory has labeled this thiol with a fluorophore to detect the binding of a variety of metals by fluorescence energy transfer or fluorescence quenching. These are steps toward the development of CA biosensors that can detect several metals.

*Phage display:* In order to display CA on the surface of filamentous phage as a functional CA-gene 3 protein (g3p) fusion protein the CA gene has been subcloned into a commercial phagemid vector, pCANTAB, which inserts the CA sequence between the g3p leader sequence and coding region. We successfully displayed active CA on the surface of filamentous phage, developed a sulfonamide affinity selection method capable of enriching CA-phage 1000-fold and used this method to screen for CA variants with altered zinc affinity. We used cassette mutagenesis to prepare a library of CA variants with mutations in three hydrophobic residues, Phe93, Phe95 and Trp97, flanking the histidine residues and used affinity selection to define the variants with the highest zinc affinity. The selected variants that are not wild-type are enriched in the following amino acids: 93-Phe, Ile, Leu, Met; 95-Met, Ile, Leu; and 97-Val, Trp. The zinc affinity of these variants is comparable to wild-type (1-20 nM) but the dissociation rate constant decreases up to 20-fold. The selected variants differ when cobalt is substituted for zinc suggesting that we can **alter metal ion specificity** as well.

SIGNIFICANCE: We have prepared CAII variants which significantly affect both the half-time for equilibration and the zinc dissociation constants. These will allow the zinc biosensor to be utilized in an array fashion for continuous monitoring. We have prepared single cysteine mutants of CA that can be specifically labeled with fluorophores for reagentless sensing of metal ions. Furthermore, we have set up a method using phage display to screen for useful CA variants.

PATENT INFORMATION: None

AWARD INFORMATION: None

PUBLICATIONS AND ABSTRACTS (for total period of grant):

1. Christianson, D.W. and Fierke, C.A. (1994) Molecular Biosensors: The Roles of Molecular and Structural Biology, Naval Res. Rev. XLVI, 30-39.

2. Kiefer, L.L., Ippolito, J.A., Christianson, D.W. and Fierke, C.A. (1994) Engineering the zinc binding site of human carbonic anhydrase II. Abstract presented 23 May 1994, Am. Soc. Biochem. Mol. Biol., The FASEB Journal 8: A1364.
3. Kiefer, L.L. and Fierke, C.A. (1994) Functional Characterization of Human Carbonic Anhydrase II Variants with Altered Zinc Binding Sites, *Biochemistry* 33,15233.
4. Thompson, R. B., Ge, Z., Patchan, M. W. and Fierke, C.A. (1995) Energy Transfer-based Fiber Optic Metal Ion Biosensor in *Advances in Fluor. Sensing Technology II: Proc. Soc. Photoopt. Instrum. Engrs. (SPIE)* vol. 2388, (J. R. Lakowicz, ed.), 138.
5. Kiefer, L. L., Paterno, S. A. and Fierke, C. A. (1995) Second Shell Hydrogen Bonds to Histidine Ligands Enhance Zinc Affinity and Catalytic Efficiency, *J. Am. Chem. Soc.* 117, 6831.
6. Ippolito, J. A., Baird, T. T., McGee, S. A., Christianson, D. W. and Fierke, C. A. (1995) Design of a Femtomolar Zinc Binding Site in Carbonic Anhydrase, *Proc. Natl. Acad. Sci.* 92, 5017.
7. Thompson, R. B., Ge, Zhengfeng, Kiefer, L. L. and Fierke, C. A. (1995) Site-Specific Mutants of Carbonic Anhydrase for Fluorescence Energy Transfer-Based Metal Ion Biosensing, *Proc. Europto Conference on Chem., Biochem. and Environ. Fibre Sensors VII*, vol. 2508, (A. V. Scheggi, ed.), Munich, FRG, SPIE, 136.
8. Huang, C.-c., Lesburg, C. A., Kiefer, L. L., Fierke, C. A. and Christianson, D. W. (1996) Reversal of the Hydrogen Bond to Zinc Ligand Histidine-119 Dramatically Diminishes Catalysis and Enhances Metal Equilibration Kinetics in Carbonic Anhydrase II, *Biochemistry* 35, 3439.
9. Thompson, R.B., Ge, Z., Patchan, M. W. and Fierke, C. A. (1996) Performance Enhancement of Fluorescence Energy Transfer-based Biosensors by Site-Directed Mutagenesis of the Transducer, *J. Biomed. Opt.* 1, 131.
10. Thompson, R. B., Ge, Z., Patchan, M., Huang, C.-C. and Fierke, C. A. (1996) Fiber Optic Biosensor for Co(II) and Cu(II) Based on Fluorescence Energy Transfer with an Enzyme Transducer, *Biosensors and Bioelectronics* 11, 557.
11. Thompson, R. B., Ge, Z., Patchan, M., Fierke, C. A., McCall, K. A., Elbaum, D. and Christianson, D. W. (1996) Determination of Multiple Analytes using a Fiber Optic Biosensor Based on Fluorescence Energy Transfer, *Progress in Biomedical Optics: Proceedings of Ultrasensitive Biochemical Diagnostics* vol. 2680, (A. Katzir, ed.), SPIE, 47.